

Title	Synthesis and Characterization of N-Heterocyclic Carbene-Coordinated Silicon Compounds Bearing a Fused-Ring Bulky Eind Group
Author(s)	Hayakawa, Naoki; Sadamori, Kazuya; Mizutani, Shinsuke; Agou, Tomohiro; Sugahara, Tomohiro; Sasamori, Takahiro; Tokitoh, Norihiro; Hashizume, Daisuke; Matsuo, Tsukasa
Citation	Inorganics (2018), 6(1)
Issue Date	2018-2-23
URL	http://hdl.handle.net/2433/233955
Right	© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This is an open access article distributed under the Creative Commons Attribution License which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited. (CC BY 4.0).
Type	Journal Article
Textversion	publisher



Article

Synthesis and Characterization of *N*-Heterocyclic Carbene-Coordinated Silicon Compounds Bearing a Fused-Ring Bulky Eind Group

Naoki Hayakawa ¹, Kazuya Sadamori ¹, Shinsuke Mizutani ¹, Tomohiro Agou ^{2,3} , Tomohiro Sugahara ² , Takahiro Sasamori ^{2,4}, Norihiro Tokitoh ², Daisuke Hashizume ⁵ and Tsukasa Matsuo ^{1,*}

¹ Department of Applied Chemistry, Faculty of Science and Engineering, Kindai University 3-4-1 Kowakae, Higashi-Osaka, Osaka 577-8502, Japan; 1544320202x@kindai.ac.jp (N.H.); 1733320213x@kindai.ac.jp (K.S.); ihsnusek555@yahoo.co.jp (S.M.)

² Institute for Chemical Research, Kyoto University, Gokasho, Uji, Kyoto 611-0011, Japan; tomohiro.agou.mountain@vc.ibaraki.ac.jp (T.A.); sugahara@boc.kuicr.kyoto-u.ac.jp (T.Su.); sasamori@nsc.nagoya-cu.ac.jp (T.Sa.); tokitoh@boc.kuicr.kyoto-u.ac.jp (N.T.)

³ Department of Biomolecular Functional Engineering, College of Engineering, Ibaraki University, 4-12-1 Nakanarusawa, Hitachi, Ibaraki 316-8511, Japan

⁴ Graduate School of Natural Sciences, Nagoya City University Yamanohata 1, Mizuho-cho, Mizuho-ku, Nagoya, Aichi 467-8501, Japan

⁵ RIKEN Center for Emergent Matter Science (CEMS), 2-1 Hirosawa, Wako, Saitama 351-0198, Japan; hashi@riken.jp

* Correspondence: t-matsuo@apch.kindai.ac.jp; Tel: +81-6-4307-3462

Received: 23 December 2017; Accepted: 14 February 2018; Published: 23 February 2018

Abstract: The reactions of the fused-ring bulky Eind-substituted 1,2-dibromodisilene, (Eind)BrSi=SiBr(Eind) (**1a**) (Eind = 1,1,3,3,5,5,7,7-octaethyl-*s*-hydrindacen-4-yl (**a**)), with *N*-heterocyclic carbenes (NHCs) (Im-Me₄ = 1,3,4,5-tetramethylimidazol-2-ylidene and Im-^{*i*}Pr₂Me₂ = 1,3-diisopropyl-4,5-dimethylimidazol-2-ylidene) are reported. While the reaction of **1a** with the sterically more demanding Im-^{*i*}Pr₂Me₂ led to the formation of the mono-NHC adduct of arylbromosilylene, (Im-^{*i*}Pr₂Me₂)→SiBr(Eind) (**2a'**), a similar reaction using the less bulky Im-Me₄ affords the bis-NHC adduct of formal arylsilyliumylidene cation, [(Im-Me₄)₂→Si(Eind)]⁺[Br[−]] (**3a**). The NHC adducts **2a'** and **3a** can also be prepared by the dehydrobromination of Eind-substituted dibromohydrosilane, (Eind)SiHBr₂ (**4a**), with NHCs. The NHC-coordinated silicon compounds have been characterized by spectroscopic methods. The molecular structures of bis-NHC adduct, [(Im-^{*i*}Pr₂Me₂)₂→Si(Eind)]⁺[Br[−]] (**3a'**), and **4a** have been determined by X-ray crystallography.

Keywords: silicon; *N*-heterocyclic carbenes; bromosilylenes; silyliumylidenes; dehydrobromination

1. Introduction

Over many years, a number of unsaturated silicon compounds have been successfully obtained by virtue of the complexation of metal ions and/or coordination of ligands (mainly Lewis bases) in addition to steric protection with bulky substituents [1–11]. Among them, the coordination chemistry of highly-reactive halosilylenes, i.e., halogen-substituted divalent Si(II) species, have attracted a lot of attention as potentially useful precursors for the construction of a wide range of silicon-containing compounds [12–17]. Figure 1 shows recent examples of coordination-stabilized arylhalosilylenes and their derivatives [18–24]. In 2010, Filippou's group reported the first *N*-heterocyclic carbene (NHC) adducts of arylchlorosilylenes bearing sterically large *m*-terphenyl groups, (Im-Me₄)→SiCl(Ar) (**1c** and **1d**) (Ar = 2,6-(Mes)₂C₆H₃ (Mes = 2,4,6-Me₃C₆H₂) (**c**)

and 2,6-(Tip)₂C₆H₃ (Tip = 2,4,6-*i*Pr₃C₆H₂) (**d**)), which were prepared by the dehydrochlorination of the aryldichlorohydrosilanes, (Ar)SiHCl₂, with NHC (Im-Me₄ = 1,3,4,5-tetramethylimidazol-2-ylidene) along with the formation of imidazolium chloride, [(Im-Me₄)H]⁺[Cl][−] [18]. The NHC adduct **Id** reacted with [Li⁺][CpMo(CO)₃][−] to afford the silylidene complex, Cp(CO)₂Mo=Si(Ar)(Im-Me₄) (**IId**), and the subsequent treatment with B(C₆H₄-4-Me)₃ produced the first silylidyne complex, Cp(CO)₂Mo≡Si(Ar) (**IIId**), featuring a metal-silicon triple bond [19,25].




Figure 1. Examples of coordination-stabilized arylhalosilylenes and their derivatives. Each one of the possible canonical forms is depicted.

In 2011, we reported on the 4-pyrrolidinopyridine (PPy) adducts of arylbromosilylenes with fused-ring bulky Rind groups, PPy→SiBr(Rind) (**IVa** and **IVb**) (Rind = 1,1,3,3,5,5,7,7-octa-R-substituted *s*-hydrindacen-4-yl; Eind (**a**: R¹ = R² = Et) and EMind (**b**: R¹ = Et, R² = Me)) [26,27], which were formed by the addition of PPy to 1,2-dibromodisilenes, (Rind)BrSi=SiBr(Rind) (**1a** and **1b**) [20]. Also in 2011, Cui reported on the related NHC-coordinated aminochlorosilylene [28], and recently Driess's group reported on the aminochlorosilylene-nickel complex [29]. In 2012, a platinum complex of arylbromosilylene, (Bbt)BrSi=Pt(PCy₃)₂ (**Ve**), was synthesized by the treatment of 1,2-dibromodisilene, (Bbt)BrSi=SiBr(Bbt) (**1e**), bearing the bulky Bbt groups (Bbt = 2,6-{CH(SiMe₃)₂}₂-4-C(SiMe₃)₃-C₆H₂) (**e**) [30] with Pt(PCy₃)₂ [21,31]. In this context, some arylbromosilylidene and arylsilylidyne complexes of nickel and platinum were reported [32–34]. In 2013, Filippou's group reported the unprecedented dicationic NHC complexes of silicon(II) and NHC adducts of iodosilyliumylidene cation Si^{II} [35]. Subsequently, we reported the reaction of 1,2-dibromodisilenes (**1b**, **1e**, and **1f**) having EMind, Bbt, and Tbb groups (Tbb = 2,6-{CH(SiMe₃)₂}₂-4-^tBu-C₆H₂) (**f**) with NHCs (Im-Me₄ and Im-ⁱPr₂Me₂) (Im-ⁱPr₂Me₂ =


1,3-diisopropyl-4,5-dimethylimidazol-2-ylidene) leading to the formation of mono-NHC adducts of arylbromosilylenes, $\text{NHC} \rightarrow \text{SiBr}(\text{Ar})$ (**VIb'** and **VIe**), and the bromide salts of the bis-NHC adducts of formal arylsilyliumylidene cations, $[(\text{NHC})_2 \rightarrow \text{Si}(\text{Ar})]^+ [\text{Br}^-]$ (**VIIf**, **VIIf'**, and **VIIf**) [22]. Inoue's group also reported the synthesis of chloride salts of bis-NHC adducts of formal arylsilyliumylidene cations, $[(\text{Im-Me}_4)_2 \rightarrow \text{Si}(\text{Ar})]^+ [\text{Cl}^-]$ (**VIIIc** and **VIIIg**) ($\text{Ar} = \text{Tip}$ (**g**)) [23] and their unique conversion to silicon analogues of acylium ions, $[(\text{Im-Me}_4)_2 \rightarrow \text{Si}(\text{O})(\text{Ar})]^+ [\text{Cl}^-]$ (**IXc** and **IXg**) [24]. Recently, Inoue's group reported on the chalcogen-atom transfer and exchange reactions of NHC-bound heavier silaacylium ions, $[(\text{Im-Me}_4)_2 \rightarrow \text{Si}(\text{E})(\text{Ar})]^+ [\text{Cl}^-]$ ($\text{E} = \text{S}, \text{Se}, \text{and Te}$) [36].

In this article, we describe the preparation and characterization of NHC-coordinated silicon compounds bearing the bulky Eind group, which have been obtained by two different synthetic procedures, i.e., the NHC-induced fragmentation of the Eind-based 1,2-dibromo-disilene and the dehydrobromination of the Eind-based dibromohydrosilane with NHCs.

2. Results and Discussions

2.1. Reactions of $(\text{Eind})\text{BrSi}=\text{SiBr}(\text{Eind})$ (**1a**) with NHCs

We first performed an NMR tube scale reaction of Eind-based 1,2-dibromodisilene, $(\text{Eind})\text{BrSi}=\text{SiBr}(\text{Eind})$ (**1a**) [20], in C_6D_6 with two equivalents of the sterically more bulky NHC, $\text{Im-}^i\text{Pr}_2\text{Me}_2$, relative to Im-Me_4 . The progress of the reaction was monitored by ^1H NMR spectroscopy, indicating the selective formation of the mono-NHC adduct of the arylbromosilylene, $(\text{Im-}^i\text{Pr}_2\text{Me}_2) \rightarrow \text{SiBr}(\text{Eind})$ (**2a'**), after overnight heating at 70°C . In the ^{29}Si NMR spectrum, only one signal was observed at $\delta = 18.0$ ppm, which is comparable to those of $(\text{Im-}^i\text{Pr}_2\text{Me}_2) \rightarrow \text{SiBr}(\text{EMind})$ (**VIb'**) ($\delta = 13.1$ ppm) and $(\text{Im-Me}_4) \rightarrow \text{SiBr}(\text{Bbt})$ (**VIe**) ($\delta = 10.9$ ppm) [22]. The ^{13}C signal at $\delta = 170.6$ ppm for **2a'** is characteristic of a carbene carbon atom, similar to those for **VIb'** ($\delta = 169.7$ ppm) and **VIe** ($\delta = 167.5$ ppm) [22]. Based on the NMR tube experiment, the mono-NHC adduct **2a'** was synthesized as an orange solid in 88% crude yield (Scheme 1).



Scheme 1. Reactions of **1a** with *N*-heterocyclic carbenes (NHCs).

We also examined the reaction of **1a** with four equivalents of $\text{Im-}^i\text{Pr}_2\text{Me}_2$. After 1-day heating at 70°C in C_6D_6 , the ^{29}Si NMR spectrum indicated the formation of a mixture containing the mono-NHC adduct **2a'** ($\delta = 18.0$ ppm) as the major product and the bis-NHC adduct of the arylsilyliumylidene cation, $[(\text{Im-}^i\text{Pr}_2\text{Me}_2)_2 \rightarrow \text{Si}(\text{Eind})]^+ [\text{Br}^-]$ (**3a'**) ($\delta = -59.6$ ppm), as the minor product. The latter ^{29}Si signal was shifted upfield compared to the former, and was similar to those of the formal

arylsilyliumylidene cations, $[(\text{NHC})_2 \rightarrow \text{Si}(\text{Eind})]^+[\text{Br}^-]$ (**VIIb** and **VIIb'**) ($\delta = -60.8$ and -75.9 ppm) and $[(\text{Im-Me}_4)_2 \rightarrow \text{Si}(\text{Tbb})]^+[\text{Br}^-]$ (**VIIIf**) ($\delta = -70.9$ ppm) [22]. However, we found that the reaction was not completed even after prolonged heating (longer than 1 week), probably due to the severe steric repulsion between the Eind group and $\text{Im-}^i\text{Pr}_2\text{Me}_2$ molecules. Thus, we were unable to isolate **3a'**. Nevertheless, single red crystals of **3a'** could be obtained from the reaction mixture, whose structure was determined by X-ray crystallography (Figure 2).




Figure 2. Molecular structure of **3a'**. The thermal ellipsoids are shown at the 50% probability level. All hydrogen atoms and benzene molecule are omitted for clarity.


Figure 2 shows a separated ion pair of **3a'** in the crystal. The closest $\text{Si} \cdots \text{Br}$ distance ($7.6669(6)$ Å) is analogous to that of **VIIb'** ($7.732(3)$ Å) [22], thus being much longer than the sum of the van der Waals radii of Si and Br (3.95 Å). The Si atom is three-coordinate adopting a distorted pyramidal geometry, which can be explained by the presence of a lone pair of electrons. The sum of the surrounding angles around the Si atom ($\Sigma\text{Si} = 327.3^\circ$) is almost the same as that of **VIIb'** ($\Sigma\text{Si} = 327.0^\circ$) [22]. The Si–C(Rind) bond length in **3a'** ($\text{Si1-C1} = 1.9482(19)$ Å) is similar to that in **VIIb'** [$1.927(8)$ Å] [22] and longer than typical Si–C bonds (ca. 1.88 Å), suggesting the high s-character of the lone pair of electrons on the Si atom and the high p-character of the Si–C(Rind) bond. The Si–C(NHC) coordination distances in **3a'** ($\text{Si1-C29} = 1.953(2)$ and $\text{Si1-C40} = 1.942(2)$ Å) are comparable to those observed in **VIIb'** ($1.955(9)$ and $1.979(8)$ Å) [22].

We next investigated the reaction of **1a** with two equivalents of the less bulky NHC, Im-Me₄, and C₆D₆. After 1 day at room temperature, two signals mainly appeared at $\delta = 73.3$ and -63.3 ppm in the ^{29}Si NMR spectrum, corresponding to the unreacted **1a** and the bis-NHC adduct of the arylsilyliumylidene cation, $[(\text{Im-Me}_4)_2 \rightarrow \text{Si}(\text{Eind})]^+[\text{Br}^-]$ (**3a**). This indicated that the NHC-aryl bromosilylene adduct, $(\text{Im-Me}_4) \rightarrow \text{SiBr}(\text{Eind})$ (**2a**), which serves as a potential intermediate, is more reactive toward Im-Me₄ compared to **1a**. When the dibromodisilene **1a** was treated with four equivalents of Im-Me₄ in benzene, the bis-NHC adduct **3a** was efficiently formed (Scheme 1). We obtained **3a** as an orange powder in 54% crude yield. The upfield-shifted ^{29}Si resonance for **3a** ($\delta = -63.3$ ppm) suggests the contribution of the canonical form due to the bis(imidazolium) adduct of a silyl anion, whose electronic structure was previously supported by the theoretical calculations of **VIIb'** and **VIIIf** [22]. In the ^{13}C NMR spectrum of **3a** in CD₃CN, one NHC carbene signal was observed at $\delta = 162.0$ ppm, comparable to those for **VIIb** ($\delta = 160.5$ ppm), **VIIb'** ($\delta = 162.4$ ppm), and **VIIIf** ($\delta = 160.4$ ppm) [22].

2.2. Reactions of (Eind)SiHBr₂ (**4a**) with NHCs

We also examined another synthetic route for the NHC-coordinated silicon compounds, i.e., the dehydrobromination of the Eind-substituted dibromohydrosilane, (Eind)SiHBr₂ (**4a**), with NHCs

(Scheme 2). The precursor (**4a**) was prepared as pale brown crystals by the dibromination of the Eind-based trihydrosilane, (Eind)SiH₃ [37,38], with allyl bromide in the presence of a catalytic amount of PdCl₂ [39]. We found that this reaction exclusively afforded **4a** even using an excess amount of allyl bromide with prolonged heating (longer than 1 week), most likely due to the steric bulkiness of the Eind group. In this context, Kunai, Ohshita, and their co-workers previously reported the selective dibromination of trihydrosilanes with CuBr₂ in the presence of CuI [40]. The formation of **4a** was deduced on the basis of the spectroscopic data (Figures S1–S4). In the ¹H NMR spectrum, the Si–H signal was found at δ = 6.89 ppm with satellite signals, due to the ²⁹Si nuclei [¹J(²⁹Si–¹H) = 288 Hz]. The ²⁹Si NMR signal appeared at δ = –28.7 ppm, similar to that of (Bbt)SiHBr₂ (δ = –28.47 ppm) [41]. The infrared spectrum exhibited a Si–H stretching band at 2317 cm^{–1} in the KBr-pellet (Figure S4) and at 2298 cm^{–1} in THF [42,43]. The molecular structure of **4a** was determined by single-crystal X-ray diffraction analysis (Figure 3). The hydrogen atom on the silicon atom was located on difference Fourier maps and isotropically refined. In the crystal, the SiHBr₂ group is fixed in one conformation with respect to the rotamer around the Si–C bond. A similar conformation was also observed in the crystal of (Eind)PCl₂ [44]. The Si–C bond length for **4a** (1.8746(18) Å) is comparable to those of typical Si–C bonds (ca. 1.88 Å).



Scheme 2. Reactions of **4a** with NHCs.




Figure 3. Molecular structure of **4a**: Side view (left), front view (right). The thermal ellipsoids are shown at the 50% probability level. The hydrogen atoms, except for the Si–H group, are omitted for clarity.

As shown in Scheme 2, the reaction of **4a** with two equivalents of Im-ⁱPr₂Me₂ proceeded more smoothly at room temperature in comparison to the reaction of **1a** with Im-ⁱPr₂Me₂ (Scheme 1), producing the mono-NHC adduct **2a'** in 59% crude yield. The reaction of **4a** with three equivalents of Im-Me₄ also afforded the bis-NHC adduct **3a** on the basis of the NMR data. In these reactions, it is essential to remove the byproducts, imidazolium bromides, [(NHC)H]⁺[Br[−]], for the isolation procedure of the silicon products, which may be considered as a disadvantage when compared to the no-byproduct strategy of using **1a** as a precursor (vide supra). Actually, the separation of **3a** and [(Im-Me₄)H]⁺[Br[−]] was found to be difficult in our experiments. However, dibromodisilene **1a** can only be obtained by a two-step synthesis from the trihydrosilane, (Eind)SiH₃; thus the bromination of (Eind)SiH₃ with *N*-bromosuccinimide (NBS) first affords the tribromosilane, (Eind)SiBr₃, then the reduction of (Eind)SiBr₃ with two equivalents of lithium naphthalenide (LiNaph) produces **1a** [20]. Therefore, the dehydrobromination of **4a** with NHCs can be considered as a convenient short-step synthesis for NHC-coordinated silylene derivatives.

3. Materials and Methods

3.1. General Procedures

All manipulations of the air- and/or moisture-sensitive compounds were performed either using standard Schlenk-line techniques or in a glove box under an inert atmosphere of argon. Anhydrous hexane, benzene, and toluene were dried by passage through columns of activated alumina and supported copper catalyst supplied by Nikko Hansen & Co., Ltd. (Osaka, Japan). Anhydrous pentane and acetonitrile were purchased from Wako Pure Chemical Industries, Ltd. (Osaka, Japan), and used without further purification. Deuterated benzene (C₆D₆, benzene-*d*₆) was dried and degassed over a potassium mirror in vacuo prior to use. Deuterated acetonitrile (CD₃CN, acetonitrile-*d*₃) was dried and distilled over calcium hydride (CaH₂) prior to use. (Eind)SiH₃ [37,38], (Eind)BrSi=SiBr(Eind) (**1a**) [20], 1,3,4,5-tetramethylimidazol-2-ylidene (Im-Me₄) [45] and 1,3-diisopropyl-4,5-dimethylimidazol-2-ylidene(Im-ⁱPr₂Me₂) [45] were prepared by the literature procedures. All other chemicals and gases were used as received.

Nuclear magnetic resonance (NMR) measurements were carried out using a JEOL ECS-400 spectrometer (399.8 MHz for ¹H, 100.5 MHz for ¹³C, and 79.4 MHz for ²⁹Si) or JEOL JNM AL-300 spectrometer (300 MHz for ¹H, 75 MHz for ¹³C, and 59 MHz for ²⁹Si) (JEOL Ltd., Tokyo, Japan). Chemical shifts (δ) are given by definition as dimensionless numbers and relative to ¹H chemical shifts of the solvents for ¹H (residual C₆D₅H in C₆D₆, ¹H(δ) = 7.15, residual CD₂HCN in CD₃CN, ¹H(δ) = 1.94), and ¹³C chemical shifts of the solvent for ¹³C (C₆D₆: ¹³C(δ) = 128.06 and CD₃CN: ¹³C(δ) = 118.26). The signal of tetramethylsilane (²⁹Si(δ) = 0.0) was used as an external standard in the ²⁹Si NMR spectra. The absolute values of the coupling constants are given in Hertz (Hz) regardless of their signs. Multiplicities are abbreviated as singlet (s), doublet (d), triplet (t), quartet (q), multiplet (m), and broad (br). The mass spectra were recorded by a JEOL JMS-T100LC AccuTOF LC-plus 4G mass spectrometer (ESI-MS) with a DART source. The elemental analyses were performed in the Microanalytical Laboratory at the Institute for Chemical Research (Kyoto University, Uji, Japan). Melting points (m.p.) were determined by a Stanford Research Systems OptiMelt instrument. We were unable to obtain a satisfactory elemental analysis for **2a'** and **3a**, probably due to their extremely high air- and moisture-sensitivity as well as a contamination of NHCs and unidentified compounds associated with some thermal decomposition (Figures S5–S10).

3.1.1. Synthesis of (Eind)SiHBr₂ (**4a**)

To a solution of (Eind)SiH₃ (4.09 g, 9.91 mmol) in toluene (30 mL) was added PdCl₂ (38.0 mg, 0.21 mmol) and allyl bromide (4.2 mL, 48.5 mmol). The reaction mixture was heated at 80 °C for 8 days. After the solvent was removed in vacuo, the residue was dissolved in hexane and the resulting mixture was centrifuged to remove the insoluble materials. The supernatant was concentrated to dryness and

the resulting residue was recrystallized from pentane to afford **4a** as pale brown crystals in 81% yield (4.58 g, 8.02 mmol).

^1H NMR (399.8 MHz, C_6D_6 , 30 °C): δ = 0.78 (t, J = 7.3 Hz, 12 H, CH_2CH_3), 0.80 (br. s, 12 H, CH_2CH_3), 1.48–1.65 (m, 8 H, CH_2CH_3), 1.76 (s, 4 H, CH_2), 2.11 (br. s, 8 H, CH_2CH_3), 6.89 (s, 1 H, satellite, $J_{\text{Si-H}}$ = 288 Hz, SiH), 7.01 (s, 1 H, ArH). ^{13}C NMR (100.5 MHz, C_6D_6 , 25 °C): δ = 9.3, 10.5 (br), 33.5, 34.2 (br), 42.6 (br \times 1), 44.8 (br \times 1), 47.9 (br \times 2), 54.7 (\times 2), 123.6, 125.1, 150.8 (one aromatic peak is broadened at 155–158); ^{29}Si NMR (79.4 MHz, C_6D_6 , 30 °C): δ = –28.7 (d, $J_{\text{Si-H}}$ = 288 Hz). IR (KBr, cm^{-1}): ν = 2317 (Si–H); IR (THF, cm^{-1}): ν = 2298 (Si–H). DART-HRMS (positive-mode) Calcd. for $\text{C}_{28}\text{H}_{46}\text{Br}_2\text{Si}$ + H: 569.1814. Found: 569.1820. Anal. Calcd. for $\text{C}_{28}\text{H}_{46}\text{Br}_2\text{Si}$: C, 58.94; H, 8.13. Found: C, 59.41; H, 8.19. Melting point (argon atmosphere in a sealed tube) 102–105 °C.

3.1.2. Synthesis of $(\text{Im-}^i\text{Pr}_2\text{Me}_2) \rightarrow \text{SiBr}(\text{Eind})$ (**2a'**)

(Method A) Reaction of $(\text{Eind})\text{BrSi}=\text{SiBr}(\text{Eind})$ (**1a**) with $\text{Im-}^i\text{Pr}_2\text{Me}_2$

A mixture of **1a** (158 mg, 0.16 mmol) and $\text{Im-}^i\text{Pr}_2\text{Me}_2$ (63.0 mg, 0.35 mmol) was dissolved in benzene (5 mL). The reaction mixture was heated overnight at 70 °C. After the solvent was removed in vacuo, the residue was washed with pentane to afford **2a'** as an orange solid in 88% crude yield (190 mg, 0.28 mmol). We were unable to isolate **2a'** in pure form, because **2a'** was not thermally stable in solution, gradually giving $\text{Im-}^i\text{Pr}_2\text{Me}_2$ and unidentified compounds (Figure S5).

^1H NMR (399.8 MHz, C_6D_6 , 60 °C): δ = 0.81–1.00 (m, 24 H, CH_2CH_3), 1.12 (br. s, 6 H, $\text{CH}(\text{CH}_3)_2-(\text{Im-}^i\text{Pr}_2\text{Me}_2)$), 1.19 (d, J = 7.0 Hz, 6 H, $\text{CH}(\text{CH}_3)_2-(\text{Im-}^i\text{Pr}_2\text{Me}_2)$), 1.61 (s, 6 H, $\text{CH}_3-(\text{Im-}^i\text{Pr}_2\text{Me}_2)$), 1.62–1.95 (m, 20 H, $\text{CH}_2 + \text{CH}_2\text{CH}_3$), 5.00–5.27 (m, 2 H, $\text{CH}(\text{CH}_3)_2-(\text{Im-}^i\text{Pr}_2\text{Me}_2)$), 6.71 (s, 1 H, ArH). ^{13}C NMR (100.5 MHz, C_6D_6 , 70 °C): δ = 9.3, 9.4, 9.9, 10.0, 10.5, 20.7, 21.4, 24.6, 33.7 (br, overlapped, $\text{Im-}^i\text{Pr}_2\text{Me}_2$ and CH_2CH_3), 42.8, 48.4, 51.1 ($\text{Im-}^i\text{Pr}_2\text{Me}_2$), 54.5, 119.9, 125.9 ($\text{Im-}^i\text{Pr}_2\text{Me}_2$), 147.2, 148.5, 153.7, 170.6 ($\text{Im-}^i\text{Pr}_2\text{Me}_2$); ^{29}Si NMR (79.4 MHz, C_6D_6 , 25 °C): δ = 18.0. HRMS (ESI, positive) Calcd. for $\text{C}_{39}\text{H}_{65}\text{BrN}_2\text{Si}$ + H: 669.4179. Found: 669.4211. Melting point (argon atmosphere in a sealed tube) 152–156 °C (dec.).

(Method B) Reaction of $(\text{Eind})\text{SiHBr}_2$ (**4a**) with $\text{Im-}^i\text{Pr}_2\text{Me}_2$

A mixture of **4a** (476 mg, 0.97 mmol) and $\text{Im-}^i\text{Pr}_2\text{Me}_2$ (352 mg, 1.95 mmol) was dissolved in benzene (7 mL). After stirring overnight at room temperature, the resulting orange suspension was filtered through a polytetrafluoroethylene (PTFE) syringe filter to remove the insoluble materials. The filtrate was concentrated to dryness and the resulting residue was washed with pentane to afford **2a'** as an orange solid in 59% crude yield (197 mg, 0.29 mmol).

3.1.3. Synthesis of $[(\text{Im-Me}_4)_2 \rightarrow \text{Si}(\text{Eind})]^+[\text{Br}]^-$ (**3a**)

(Method A) Reaction of $(\text{Eind})\text{BrSi}=\text{SiBr}(\text{Eind})$ (**1a**) with Im-Me_4

A mixture of **1a** (102 mg, 0.11 mmol) and Im-Me_4 (54 mg, 0.43 mmol) was dissolved in benzene (6 mL). After stirring for 1 day at room temperature, the resulting orange solid was separated and washed with a mixture of hexane and benzene to afford **3a** as an orange powder in 54% crude yield (85.2 mg, 0.12 mmol). We were unable to isolate **3a** in pure form, because **3a** was not thermally stable in solution leading to the contamination of Im-Me_4 and unidentified compounds (Figure S8).

^1H NMR (399.8 MHz, CD_3CN , 20 °C): δ = 0.61 (br. t, J = 6.3 Hz, 12 H, CH_2CH_3), 0.80 (t, J = 7.3 Hz, 12 H, CH_2CH_3), 1.50–1.70 (m, 8 H, CH_2CH_3), 1.75 (br. s, 4 H, CH_2), 1.93–1.97 (m, overlapped, 8 H, CH_2CH_3), 2.15 (s, 12 H, $\text{CH}_3-(\text{Im-Me}_4)$), 3.25 (br. s, 12 H, $\text{CH}_3-(\text{Im-Me}_4)$), 6.75 (s, 1 H, ArH). ^{13}C NMR (100.5 MHz, CD_3CN , 19 °C): δ = 9.2, 9.4, 10.0 (br, Im-Me_4), 34.1 (br, overlapped, Im-Me_4 and CH_2CH_3), 42.8, 48.1, 54.2, 121.8, 128.5 (Im-Me_4), 136.5, 150.7, 162.0 (Im-Me_4) (one aromatic peak is overlapped). ^{29}Si NMR

(79.4 MHz, CD₃CN, 25 °C): $\delta = -63.3$. DART-HRMS (positive-mode) Calcd. for C₄₂H₆₉BrN₄Si + H: 737.4553. Found: 737.4562. Melting point (argon atmosphere in a sealed tube) 169–174 °C (dec.).

(Method B) Reaction of (Eind)SiHBr₂ (**4a**) with Im-Me₄

A mixture of **4a** (70.3 mg, 0.12 mmol) and Im-Me₄ (50.6 mg, 0.41 mmol) was dissolved in benzene (6 mL). After stirring for 1 day at room temperature, an orange suspension was formed. An insoluble orange solid was collected by filtration, whose ¹H NMR spectrum indicated the formation of a mixture of **4a** and [(Im-Me₄)H]⁺[Br[−]].

3.2. X-ray Crystallographic Studies of **3a'** and **4a**

Single crystals suitable for X-ray diffraction measurements were obtained from benzene for **3a'** and from hexane for **4a**. Intensity data were collected using a Rigaku XtaLAB P200 with a PILATUS 200K detector for **3a'** and a Rigaku AFC-8 with a Saturn 70 CCD detector for **4a** (Rigaku Corporation, Tokyo, Japan). All measurements were carried out using Mo K α radiation ($\lambda = 0.71073$ Å). The integration and scaling of the diffraction data were carried out using the programs CrysAlisPro [46] for **3a'** and CrystalClear [47] for **4a**. Lorentz, polarization, and absorption corrections were also performed. The structures were solved by an iterative method with the program of SHELXT [48], and refined by a full-matrix least-squares method on F^2 for all the reflections using the program SHELXL-2017/1 [49]. The non-hydrogen atoms were refined by applying anisotropic temperature factors. Positions of all the hydrogen atoms were geometrically calculated, and refined as riding models. The Si–H hydrogen atom was located on difference Fourier maps and isotropically refined. Full details of the crystallographic analysis and accompanying CIF files can be obtained free of charge from the Cambridge Crystallographic Data Centre (CCDC numbers 1811699 and 1811700) via <http://www.ccdc.cam.ac.uk/conts/retrieving.html> (or from the CCDC, 12 Union Road, Cambridge CB2 1EZ, UK; Fax: +44-1223-336033; E-mail: deposit@ccdc.cam.ac.uk).

3.2.1. [(Im-ⁱPr₂Me₂)₂→Si(Eind)]⁺[Br[−]] (**3a'**)

C₅₀H₈₅BrN₄Si·C₆H₆, $M = 928.32$, crystal size $0.36 \times 0.15 \times 0.12$ mm, triclinic, space group $P-1$ (#2), $a = 10.6548(3)$ Å, $b = 12.0149(2)$ Å, $c = 21.7366(5)$ Å, $\alpha = 80.5199(18)^\circ$, $\beta = 82.343(2)^\circ$, $\gamma = 72.816(2)^\circ$, $V = 2611.54(11)$ Å³, $Z = 2$, $D_x = 1.181$ g cm^{−3}, $\mu(\text{Mo } K\alpha) = 0.849$ mm^{−1}, 65594 reflections collected, 13761 unique reflections, and 579 refined parameters. The final $R(F)$ value was 0.0491 [$I > 2\sigma(I)$]. The final $R_w(F^2)$ value was 0.1364 (all data). The goodness-of-fit on F^2 was 1.031.

3.2.2. (Eind)SiHBr₂ (**4a**)

C₂₈H₄₆Br₂Si, $M = 570.56$, crystal size $0.16 \times 0.17 \times 0.41$ mm, triclinic, space group $P-1$ (#2), $a = 7.972(3)$ Å, $b = 11.070(4)$ Å, $c = 16.621(5)$ Å, $\alpha = 89.972(4)^\circ$, $\beta = 80.770(3)^\circ$, $\gamma = 73.314(5)^\circ$, $V = 1385.1(8)$ Å³, $Z = 2$, $D_x = 1.368$ g cm^{−3}, $\mu(\text{Mo } K\alpha) = 2.992$ mm^{−1}, 22599 reflections collected, 6332 unique reflections, and 292 refined parameters. The final $R(F)$ value was 0.0298 [$I > 2\sigma(I)$]. The final $R_w(F^2)$ value was 0.0803 (all data). The goodness-of-fit on F^2 was 1.016.

4. Conclusions

We have synthesized some new NHC-coordinated silicon species having the fused-ring bulky Eind group by two methods; one is via the reactions of the stable diaryldibromodisilene, (Eind)BrSi=SiBr(Eind) (**1a**), with NHCs, and the other is the dehydrobromination of the aryldibromohydrosilane, (Eind)SiHBr₂ (**4a**), with NHCs. In both synthetic pathways, we have mainly obtained the mono-NHC adduct of the arylbromosilylene, (Im-ⁱPr₂Me₂)→SiBr(Eind) (**2a'**), and the bis-NHC adduct of the formal arylsilyliumylidenecation, [(Im-Me₄)₂→Si(Eind)]⁺[Br[−]] (**3a**), depending on the steric bulk of the NHCs (Im-ⁱPr₂Me₂ vs. Im-Me₄). Further studies on the reactivities of the NHC-coordinated silicon compounds are now in progress.

Supplementary Materials: The following are available online at www.mdpi.com/2304-6740/6/1/30/s1, S1: NMR spectra of **4a**, **2a'**, and **3a** and IR spectrum of **4a** (PDF), S2: crystallographic details for **4a** and **3a'** (CIF) and S3: cif-checked files (PDF).

Acknowledgments: This study was supported by Grants-in-Aid for Scientific Research on Innovative Areas “Stimuli-responsive Chemical Species for the Creation of Functional Molecules (No. 2408)” (JSPS KAKENHI Grant Nos. JP20109003 for Tsukasa Matsuo, JP15H00964 for Daisuke Hashizume, and JP20109013 for Norihiro Tokitoh), Scientific Research (B) (No. JP15H03788 for Tsukasa Matsuo and JP15H03777 for Takahiro Sasamori), Young Scientists (A) (No. 15H05477 for Tomohiro Agou) and Challenging Exploratory Research (No. 16K13953 for Tomohiro Agou). This study was partially supported by a MEXT-Supported Program for the Strategic Research Foundation at Private Universities 2014–2018 subsidy from MEXT and Kindai University. We thank the Collaborative Research Program of The Institute for Chemical Research, Kyoto University (grants #2016-94 and #2017-99). Naoki Hayakawa and Tomohiro Sugahara acknowledge the support by Grants-in-Aid for JSPS Fellows from JSPS (Nos. JP16J01036 and JP16J05501).

Author Contributions: Naoki Hayakawa, Kazuya Sadamori, Shinsuke Mizutani, Tomohiro Agou, and Tomohiro Sugahara performed the experiments. Naoki Hayakawa and Daisuke Hashizume carried out the X-ray crystallographic analysis. Tomohiro Sugahara, Norihiro Tokitoh, and Tsukasa Matsuo designed the experiments and co-directed the project. Naoki Hayakawa, Kazuya Sadamori, and Tsukasa Matsuo co-wrote the paper. Takahiro Sasamori, Norihiro Tokitoh, Daisuke Hashizume, and Tsukasa Matsuo reviewed and approved the final manuscript. All authors contributed to the discussions.

Conflicts of Interest: The authors declare no conflict of interest.

References

1. Kira, M.; Iwamoto, T. Progress in the chemistry of stable disilenes. *Adv. Organomet. Chem.* **2006**, *54*, 73–148.
2. Wang, Y.; Robinson, G.H. Unique homonuclear multiple bonding in main group compounds. *Chem. Commun.* **2009**, 5201–5213. [[CrossRef](#)] [[PubMed](#)]
3. Mizuhata, Y.; Sasamori, T.; Tokitoh, N. Stable heavier carbene analogues. *Chem. Rev.* **2009**, *109*, 3479–3511. [[CrossRef](#)] [[PubMed](#)]
4. Fischer, R.C.; Power, P.P. π -Bonding and lone pair effect in multiple bonds involving heavier main group elements: Developments in the new millennium. *Chem. Rev.* **2010**, *110*, 3877–3923. [[CrossRef](#)] [[PubMed](#)]
5. Lee, V.Y.; Sekiguchi, A. *Organometallic Compounds of Low-Coordinate Si, Ge, Sn and Pb: From Phantom Species to Stable Compounds*; John Wiley & Sons Ltd.: Chichester, UK, 2010; ISBN 978-0-470-72543-6.
6. Asay, M.; Jones, C.; Driess, M. *N*-Heterocyclic carbene analogues with low-valent group 13 and group 14 elements: Syntheses, structures, and reactivities of a new generation of multitasking ligands. *Chem. Rev.* **2011**, *111*, 354–396. [[CrossRef](#)] [[PubMed](#)]
7. Yao, S.; Xiong, Y.; Driess, M. Zwitterionic and donor-stabilized *N*-heterocyclic silylenes (NHSis) for metal-free activation of small molecules. *Organometallics* **2011**, *30*, 1748–1767. [[CrossRef](#)]
8. Kira, M. Bonding and structure of disilenes and related unsaturated Group-14 element compounds. *Proc. Jpn. Acad. Ser. B Phys. Biol. Sci.* **2012**, *88*, 167–191. [[CrossRef](#)] [[PubMed](#)]
9. Scheschewitz, D. *Functional Molecular Silicon Compounds II: Low Oxidation States*; Springer: Basel, Switzerland, 2014; ISBN 978-3-319-03734-9.
10. Wang, Y.; Robinson, G.H. *N*-heterocyclic carbene—Main-group chemistry: A rapidly evolving field. *Inorg. Chem.* **2014**, *53*, 11815–11832. [[CrossRef](#)] [[PubMed](#)]
11. Ghadwal, R.S.; Azhakar, R.; Roesky, H.W. Dichlorosilylene: A high temperature transient species to an indispensable building block. *Acc. Chem. Res.* **2013**, *46*, 444–456. [[CrossRef](#)] [[PubMed](#)]
12. Ghadwal, R.S.; Roesky, H.W.; Merkel, S.; Henn, J.; Stalke, D. Lewis base stabilized dichlorosilylene. *Angew. Chem. Int. Ed.* **2009**, *48*, 5683–5686. [[CrossRef](#)] [[PubMed](#)]
13. Filippou, A.C.; Chernov, O.; Schnakenburg, G. SiBr₂(Idipp): A stable *N*-heterocyclic carbene adduct of dibromosilylene. *Angew. Chem. Int. Ed.* **2009**, *48*, 5687–5690. [[CrossRef](#)] [[PubMed](#)]
14. Roy, S.; Stollberg, P.; Herbst-Irmer, R.; Stalke, D.; Andrada, D.M.; Frenking, G.; Roesky, H.W. Carbene-dichlorosilylene stabilized phosphinidenes exhibiting strong intramolecular charge transfer transition. *J. Am. Chem. Soc.* **2015**, *137*, 150–153. [[CrossRef](#)] [[PubMed](#)]
15. Hickox, H.P.; Wang, Y.; Xie, Y.; Wei, P.; Schaefer, H.F., III; Robinson, G.H. Push-pull stabilization of parent monochlorosilylenes. *J. Am. Chem. Soc.* **2016**, *138*, 9799–9802. [[CrossRef](#)] [[PubMed](#)]
16. Geiß, D.; Arz, M.I.; Straßmann, M.; Schnakenburg, G.; Filippou, A.C. Si=P double bonds: experimental and theoretical study of an NHC-stabilized phosphasilylenylidene. *Angew. Chem. Int. Ed.* **2015**, *54*, 2739–2744.

17. Ghana, P.; Arz, M.I.; Das, U.; Schnakenburg, G.; Filippou, A.C. Si=Si double bonds: Synthesis of an NHC-stabilized disilavinylidene. *Angew. Chem. Int. Ed.* **2015**, *54*, 9980–9985. [[CrossRef](#)] [[PubMed](#)]
18. Filippou, A.C.; Chernov, O.C.; Blom, B.; Stumpf, K.W.; Schnakenburg, G. Stable *N*-heterocyclic carbene adducts of arylchlorosilylenes and their germanium homologues. *Chem. Eur. J.* **2010**, *16*, 2866–2872. [[CrossRef](#)] [[PubMed](#)]
19. Filippou, A.C.; Chernov, O.C.; Stumpf, K.W.; Schnakenburg, G. Metal-silicon triple bonds: The molybdenumsilylidyne complex [Cp(CO)₂Mo≡Si-Ar]. *Angew. Chem. Int. Ed.* **2010**, *49*, 3296–3300. [[CrossRef](#)] [[PubMed](#)]
20. Suzuki, K.; Matsuo, T.; Hashizume, D.; Tamao, K. Room-temperature dissociation of 1,2-dibromodisilenes to bromosilylenes. *J. Am. Chem. Soc.* **2011**, *133*, 19710–19713. [[CrossRef](#)] [[PubMed](#)]
21. Agou, T.; Sasamori, T.; Tokitoh, N. Synthesis of an arylbromosilylene-platinum complex by using a 1,2-dibromodisilene as a silylene source. *Organometallics* **2012**, *31*, 1150–1154. [[CrossRef](#)]
22. Agou, T.; Hayakawa, N.; Sasamori, T.; Matsuo, T.; Hashizume, D.; Tokitoh, N. Reactions of diaryldibromodisilenes with *N*-heterocyclic carbenes: Formation of formal bis-NHC adducts of silyliumylidene cations. *Chem. Eur. J.* **2014**, *20*, 9246–9249. [[CrossRef](#)] [[PubMed](#)]
23. Ahmad, S.U.; Szilvási, T.; Inoue, S. A facile access to a novel NHC-stabilized silyliumylidene ion and C–H activation of phenylacetylene. *Chem. Commun.* **2014**, *50*, 12619–12622. [[CrossRef](#)] [[PubMed](#)]
24. Ahmad, S.U.; Szilvási, T.; Irran, E.; Inoue, S. An NHC-stabilized silicon analogue of acylium ion: Synthesis, structure, reactivity, and theoretical studies. *J. Am. Chem. Soc.* **2015**, *137*, 5828–5836. [[CrossRef](#)] [[PubMed](#)]
25. Mork, B.V.; Tilley, T.D. Multiple bonding between silicon and molybdenum: A transition-metal complex with considerable silylyne character. *Angew. Chem. Int. Ed.* **2003**, *42*, 357–360. [[CrossRef](#)] [[PubMed](#)]
26. Matsuo, T.; Suzuki, K.; Fukawa, T.; Li, B.L.; Ito, M.; Shoji, Y.; Otani, T.; Li, L.C.; Kobayashi, M.; Hachiya, M.; et al. Synthesis and structures of a series of bulky “Rind-Br” based on a rigid fused-ring *s*-hydrindacene skeleton. *Bull. Chem. Soc. Jpn.* **2011**, *84*, 1178–1191. [[CrossRef](#)]
27. Matsuo, T.; Tamao, K. Fused-ring bulky “Rind” groups producing new possibilities in elemento-organic chemistry. *Bull. Chem. Soc. Jpn.* **2015**, *88*, 1201–1220. [[CrossRef](#)]
28. Cui, H.; Cui, C. Silylation of *N*-heterocyclic carbene with aminochlorosilane and -disilane: Dehydrohalogenation vs. Si–Si bond cleavage. *Dalton Trans.* **2011**, *40*, 11937–11940. [[CrossRef](#)] [[PubMed](#)]
29. Hadlington, T.J.; Szilvási, T.; Driess, M. Silylene-nickel promoted cleavage of B–O bonds: From catechol borane to the hydroborylene ligand. *Angew. Chem. Int. Ed.* **2017**, *56*, 7470–7474. [[CrossRef](#)] [[PubMed](#)]
30. Sasamori, T.; Hironaka, K.; Sugiyama, Y.; Takagi, N.; Nagase, S.; Hosoi, Y.; Furukawa, Y.; Tokitoh, N. Synthesis and reactions of a stable 1,2-diaryl-1,2-dibromodisilene: A precursor for substituted disilenes and a 1,2-diaryldisilyne. *J. Am. Chem. Soc.* **2008**, *130*, 13856–13857. [[CrossRef](#)] [[PubMed](#)]
31. Liao, W.-H.; Ho, P.-Y.; Su, M.-D. Mechanisms for the reactions of Group 10 transition metal complexes with metal-group 14 element bonds, Bbt(Br)E=M(PCy₃)₂ (E = C, Si, Ge, Sn, Pb; M = Pd and Pt). *Inorg. Chem.* **2013**, *52*, 1338–1348. [[CrossRef](#)] [[PubMed](#)]
32. Filippou, A.C.; Hoffmann, D.; Schnakenburg, G. Triple bonds of niobium with silicon, germanium and tin: The tetrylidyne complexes [(κ³-tmpts)(CO)₂Nb≡E–R] (E = Si, Ge, Sn; tmpts = MeSi(CH₂OMe)₃; R = aryl). *Chem. Sci.* **2017**, *8*, 6290–6299. [[CrossRef](#)]
33. Blom, B. Reactivity of Ylenes at Late Transition Metal Centers. Dissertation, University of Bonn, Göttingen, Germany, 2011.
34. Papazoglou, I. Unprecedented Tetrylidyne Complexes of Group 6 and Group 10 Metals. Dissertation, University of Bonn, Dr. Hut Verlag, München, Germany, 28 May 2017.
35. Filippou, A.C.; Lebedev, Y.N.; Chernov, O.; Straßmann, M.; Schnakenburg, G. Silicon(II) coordination chemistry: *N*-Heterocyclic carbene complexes of Si²⁺ and SiI⁺. *Angew. Chem. Int. Ed.* **2013**, *52*, 6974–6978. [[CrossRef](#)] [[PubMed](#)]
36. Sarkar, D.; Wendel, D.; Ahmad, S.U.; Szilvási, T.; Pöthig, A.; Inoue, S. Chalcogen-atom transfer and exchange reactions of NHC-stabilized heavier silaacylium ions. *Dalton Trans.* **2017**, *46*, 16014–16018. [[CrossRef](#)] [[PubMed](#)]
37. Fukazawa, A.; Li, Y.; Yamaguchi, S.; Tsuji, H.; Tamao, K. Coplanar oligo(*p*-phenylenedisilenylene)s based on the octaethyl-substituted *s*-hydrindacenyl groups. *J. Am. Chem. Soc.* **2007**, *129*, 14164–14165. [[CrossRef](#)] [[PubMed](#)]

38. Kobayashi, M.; Hayakawa, N.; Nakabayashi, K.; Matsuo, T.; Hashizume, D.; Fueno, H.; Tanaka, K.; Tamao, K. Highly coplanar (*E*)-1,2-di(1-naphthyl)disilene involving a distinct CH- π interaction with the perpendicularly oriented protecting Eind group. *Chem. Lett.* **2014**, *43*, 432–434. [CrossRef]
39. Iwata, A.; Toyoshima, Y.; Hayashida, T.; Ochi, T.; Kunai, A.; Ohshita, J. PdCl₂ and NiCl₂-catalyzed hydrogen-halogen exchange for the convenient preparation of bromo- and iodosilanes and germanes. *J. Organomet. Chem.* **2003**, *667*, 90–95. [CrossRef]
40. Kunai, A.; Ochi, T.; Iwata, A.; Ohshita, J. Synthesis of bromohydrosilanes: Reactions of hydrosilanes with CuBr₂ in the presence of CuI. *Chem. Lett.* **2001**, 1228–1229. [CrossRef]
41. Agou, T.; Sugiyama, Y.; Sasamori, T.; Sakai, H.; Furukawa, Y.; Takagi, N.; Guo, J.-D.; Nagasa, S.; Hashizume, D.; Tokitoh, N. Synthesis of Kinetically Stabilized 1,2-Dihydrodisilenes. *J. Am. Chem. Soc.* **2012**, *134*, 4120–4123. [CrossRef] [PubMed]
42. Simons, R.S.; Haubrich, S.T.; Mork, B.V.; Niemeyer, M.; Power, P.P. The Syntheses and Characterization of the Bulky Terphenylsilanes and Chlorosilanes 2,6-Mes₂C₆H₃SiCl₃, 2,6-Trip₂C₆H₃SiCl₃, 2,6-Mes₂C₆H₃SiHCl₂, 2,6-Trip₂C₆H₃SiHCl₂, 2,6-Mes₂C₆H₃SiH₃, 2,6-Trip₂C₆H₃SiH₃ and 2,6-Mes₂C₆H₃SiCl₂SiCl₃. *Main Group Chem.* **1998**, *2*, 275–283. [CrossRef]
43. Weidemann, N.; Schnakenburg, G.; Filippou, A.C. Novel silanes with sterically demanding aryl substituents. *Z. Anorg. Allg. Chem.* **2009**, *635*, 253–259. [CrossRef]
44. Li, B.; Tsujimoto, S.; Li, Y.; Tsuji, H.; Tamao, K.; Hashizume, D.; Matsuo, T. Synthesis and characterization of diphosphenes bearing fused-ring bulky Rind groups. *Heteroat. Chem.* **2014**, *25*, 612–618. [CrossRef]
45. Kuhn, S.; Kratz, T. Synthesis of imidazol-2-ylidenes by reduction of imidazole-2(3*H*)-thiones. *Synthesis* **1993**, 561–562. [CrossRef]
46. *CrysAlisPro*; Agilent Technologies Ltd.: Yarnton, Oxfordshire, UK, 2014.
47. *CrystalClear*; Rigaku/MSK Inc.: The Woodlands, TX, USA, 2005.
48. Sheldrick, G.M. *SHELXT*—Integrated space-group and crystal-structure determination. *Acta Crystallogr. Sect. A* **2015**, *A71*, 3–8. [CrossRef] [PubMed]
49. Sheldrick, G.M. Crystal structure refinement with *SHELXL*. *Acta Crystallogr. Sect. C* **2015**, *C71*, 3–8.



© 2018 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<http://creativecommons.org/licenses/by/4.0/>).